

A service of the National Library of Medicine
and the National Institutes of Health

www.pubmed.gov

My NCBI [?]
[Sign In] [Register]

All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Books

Search PubMed for Go Clear

☒ Limits ☒ Preview/Index ☒ History ☒ Clipboard ☒ Details

Display AbstractPlus Show 20 Sort by Send to

All: 1 Review: 0

1: Arch Dermatol. 1990 Apr;126(4):487-93.

Links

Comment in:

Arch Dermatol. 1990 Nov;126(11):1516-7.

Sulfasalazine improves psoriasis. A double-blind analysis.

Gupta AK, Ellis CN, Siegel MT, Duell EA, Griffiths CE, Hamilton TA, Nickoloff BJ, Voorhees JJ.

Department of Dermatology, University of Michigan Medical Center, Ann Arbor 48109-0314.

In an 8-week double-blind trial of sulfasalazine for the treatment of moderate-to-severe psoriasis, 23 and 27 patients received the active and placebo tablets, respectively. At the end of the double-blind phase, there were 17 assessable patients receiving sulfasalazine; 7 (41%) had marked improvement, 7 (41%) had moderate improvement, and 3 (18%) demonstrated minimal change. Only 1 patient receiving placebo demonstrated moderate improvement, whereas the rest had minimal improvement or worsening of psoriasis. When the randomization code was broken, patients receiving sulfasalazine were allowed to continue therapy for an additional 4 weeks in an open manner, while those using placebo left the study. Six of 23 patients discontinued sulfasalazine therapy during the double-blind phase because of side effects, 4 due to the development of a cutaneous eruption and 2 due to nausea. Fourteen of 17 patients, who were assessable at the end of the 8-week double-blind phase, completed the additional 4 weeks of sulfasalazine therapy. Of these 14 patients, marked improvement occurred in 8 (57%), moderate improvement in 2 (14%), and minimal improvement in 4 (29%), compared with pretherapy. The low incidence of severe side effects makes sulfasalazine a consideration for oral therapy in patients whose disease severity does not justify use of methotrexate, etretinate, or psoralen plus UV-A, but whose disease severity is too widespread for safe and practical use of topical corticosteroids.

PMID: 1690970 [PubMed - indexed for MEDLINE]

Display AbstractPlus Show 20 Sort by Send to

Related Links

Sulfasalazine: a potential psoriasis therapy? [J Am Acad Dermatol. 1989]

A double-blind, placebo-controlled trial of acitretin for the treatment of psoriasis [J Am Acad Dermatol. 1989]

Topical PTH (1-34) is a novel, safe and effective treatment for psoriasis: a randomized self-controlled trial and an open trial. [Br J Dermatol. 2003]

Sulphasalazine in psoriatic arthritis: a double-blind placebo-controlled study [J Rheumatol. 1990]

Olsalazine sodium in the treatment of ulcerative colitis among patients intolerant of sulfasalazine. A prospective, randomized, placebo-controlled, double-blind, dose-ranging clinical trial. [Gastroenterology. 1987]

See all Related Articles...

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Privacy Statement | Freedom of Information Act | Disclaimer

Oct 17 2006 07:23:06